

Yellow archangel

Lamium galeobdolon

Integrated Pest Management Control Measures

Mechanical

Cutting or pulling plants alone will not control this species. Plant readily re-sprouts from any fragments left behind and pulling must be repeated several times in a season over several years.

Cultural

Patches can be covered with cardboard and 3-4 inches of mulch material or with cloth and mulch. Entire patch must be covered this way and monitored along edges and gaps for re-growth.

Biological

No biological controls are currently available in the United States.

Chemical *Always read herbicide label before applying*

Timing and products are variable; for best results please consult the chemical label for best recommendations on when to treat.



FOLIAR SPRAY			Calendar/Plant Stage
Chemical	Brand Names	Label Use	
Imazapyr	Habitat, Arsenal	Upland, non-cropland	Apply to actively growing plants. Application during this time has increased effectiveness because the herbicide moves into plant more quickly. Treating with a mix of triclopyr and glyphosate may achieve better results. Re-treatment with herbicides is likely necessary.
MCP + 2,4D + dicamba	Ortho Weed B Gon	Turf, ornamental	
triclopyr	Garlon 3A	Non-cropland	
triclopyr + 2,4-D	Crossbow	Rangeland, Permanent grass pasture, non-cropland	
metsulfuron	Escort	Pasture, Rangeland and non-cropland	
glyphosate Non Selective Herbicide	RoundUp	Non-cropland	



Cowlitz County Noxious Weed Control Board
 Administration Annex Building, 1st Floor ■ Kelso, WA 98626
 Phone: 360-577-3117 ■ Fax: 360-425-7760
 E-mail: velazqueza@co.cowlitz.wa.us