



Why is tansy ragwort a problem?

Tansy ragwort is an invasive, toxic weed. When prevalent, tansy ragwort is one of the most common causes of poisoning in cattle and horses, caused by consumption of the weed found in pasture, hay or silage. Milk produced by affected cows and goats can contain toxins.

Stock does not reject or avoid it in hay or silage and its poisonous alkaloids are unaffected by drying. Honey from tansy ragwort also contains the alkaloids. The highest risk is after the plants have been cut or when mixed in with hay, because the plants are not as bitter but just as toxic.



Recognizing tansy ragwort

Tansy ragwort is a tap rooted biennial and sometimes a perennial herb growing up to four feet tall.

It produces flowerheads that are flat topped clusters. Flowerheads are yellow with many disk flowers and 13 ray flowers. Overall, flowerheads have a daisy-like appearance and bloom June to August.

Tansy ragwort spends the first year in the rosette stage with dark green basal leaves that ap-

pear ruffled. The leaf underside is somewhat hairy, and appears whitish. During the second year, one



or several flowering stems bolt. The leaves found on the flowering stem are alternate and sessile.



which can lie dormant in the soil for as long as 15 years.

Controlling noxious weeds on your property is your responsibility and the law.

Chapter 17.10 RCW, County Code Title 7

YOU can help stop the spread of noxious weeds

- Report infestations
- Actively control noxious weeds on your property
- Contact Clark County's Vegetation Management Program for more information on species ID and recommended control methods
- Spread the word about noxious weeds, and why controlling them is so important

Remember, weeds are everyone's problem. Controlling noxious weeds on your property is your responsibility and the law.

Online Resources

Clark County Noxious Weed Program www.clark.wa.gov/weed

Washington State Noxious Weed Control Board Www.nwcb.wa.gov

Early Detection & Distribution Mapping System Mobile App: www.eddmaps.org/west

Scan this QR code to download

For more information:

Vegetation Management Program (360) 397-6140 email: weed.management@clark.wa.gov

For other formats, contact the Clark County ADA Office: Voice (360) 397-2322, **Relay** 711 or (800) 833-6388, **Fax** (360) 397-6165, **E-mail** ADA@clark.wa.gov.

Clark County Public Works Vegetation Management (360) 397-2121 www.clark.wa.gov/weed Tansy ragwort



CLARK COUNTY PUBLIC WORKS VEGETATION MANAGEMENT PROGRAM



Why control noxious weeds in Clark County?

Noxious weeds are non-native plants that can be toxic, destructive, competitive and difficult to control once established.

Economics - Noxious weeds cost the United States on average 30.6 billion dollars each year in decreased land value, money and time spent in control efforts, lower crop yields, reduced forage quality and impacts on animal health.

Environmental health - Noxious weeds displace native species, destroy natural habitat, clog waterways and increase erosion and fire risk.

Human & animal health - Many noxious weeds are toxic to humans and livestock. Contact or ingestion of some species can lead to serious health problems or death.

Recreation - Noxious weeds hurt recreation opportunities such as bird watching and fishing through reduced accessibility and destruction of native landscapes.

Integrated Weed Management (IWM)



An Integrated Weed Management plan is an ongoing, continuing cycle of weed prevention, control, monitoring, evaluation and planning

Managing weeds with Integrated Weed Management

The most effective way to manage weed infestations is to use a combination of control methods specific to the problem weed, where it is in its growth-cycle, and the location where it is growing. This approach is called integrated weed management, or IWM, which uses biological, mechanical, cultural, and chemical (herbicide) control methods that treat the problem weed yet protect human health, habitat, water, and other natural resources.

Prevention is better than control - The best control method of all is to prevent weeds in the first place. IWM starts with understanding the soil, water, natural resources and human impacts and uses on a site. For example, weeds often invade due to overgrazing, bare soil, or other factors that should be corrected for the control measures to be fully effective.

Long-term effectiveness – A good IWM plan is more effective than complete reliance on herbicide management. While not all control methods are useful for all weed species, taking an integrated approach to weed management can greatly increase the effectiveness of your efforts. As weed control is not a one-time fix, an IWM strategy should be practical, adaptable, cost-efficient, and effective.

IWM control recommendations for tansy ragwort

IWA	N			Effectiveness of control method								THE WEED CONTROL TOOL		
control		Control method		Sn	Small/backyard site				large/r	ural site				
typ	е			Good	Fair	Poor	N/A	Good	Fair	Poor	N/A		VSICAL	
		digging									•		owing	
Physical mechanic	al &	hand-pulling									•		ulling	
	nical	mowing								•		di	gging	
		tilling					•				•			
		bark mulch			•					•				
	ltural	black plastic									•			
Cultu		cover crop			•									
		native plant restoration											2750	
		soil amendment										Carl Conter		
Dielog	vical	managed grazing					•				•	CULTURAL	BIOL	
ылод	gical	weed-feeding insects										soil amendments cover crops•mulch	weed-ea	
			Product examples *									native plants	manag	
	emical	aminopyralid	Milestone									C SKEL		
Chem		glyphosate	Roundup, AquaNeat, Rodeo	•					•					
		triclopyr amine	Garlon 3A, Lilly Miller Brush Killer	•				•						

* Brand names are listed as an example only. Other commercial products may contain the listed chemical control. Clark County does not endorse any product or brand name. Always read and follow the herbicide label. For more information on specific herbicides, please contact Vegetation Management.

BOX

Integrated Weed Managment uses multiple tools in combination for the most effective weed control.

GICAL

ing insect d grazing

