



Please contact your County Noxious Weed Control Board to learn which Class C species are selected for control in your area.

Himalayan blackberry (above) is widespread in many parts of the state. Counties may require landowners to control this species or may opt to provide educational resources and encourage voluntary control.



Canada thistle (above) is widespread throughout Washington. Several counties selected this species for control in their county last year, including Adams County, Clark County, Franklin County, and Spokane County, among others.



Noxious weeds that are typically widespread in WA or are of special interest to the state's agricultural industry. The State Weed Board provides educational resources about these species but does not require control of them. The Class C status allows counties to require control if locally desired. Other counties may choose to provide education or technical consultation.

Class C Weeds

County Contacts

Adams	(509) 659-1806
Adam - District #1	(509) 488-9023
Asotin	(509) 243-2032
Benton	(509) 943-6005
Benton - District #1	(509) 786-5609
Chelan	(509) 667-6576
Clallam	(360) 417-2442
Clark	(360) 397-6140
Columbia	(509) 382-9760
Cowlitz	(360) 577-3117
Douglas (WSU)	(509) 745-8531 Ext. 6352
Ferry	(509) 775-5225 Ext. 1111
Franklin	(509) 545-3847
Garfield	(509) 843-1913
Grant	(509) 754-2011 Ext. 4710
Grant - District #1	(509) 750-8948
Grant - District #2	(509) 288-2483
Grant - District #3	(509) 398-1870
Grays Harbor	(360) 482-2265
Island	(360) 678-7992
Jefferson	(360) 379-5610 Ext. 205
King	(206) 477-9333
Kitsap	(360) 337-7157 Ext. 6261
Kittitas	(509) 962-7007
Klickitat	(509) 773-5810
Lewis	(360) 740-1215
Lincoln	(509) 725-3646 Ext. 521
Mason	(360) 427-9670 Ext. 592
Okanogan	(509) 422-7165
Pacific	(360) 875-9425
Pend Oreille	(509) 447-2402
Pierce	(253) 798-7263
San Juan	(360) 376-3499
Skagit	(360) 416-1467
Skamania	(509) 427-3941

Please contact your County Noxious Weed Control Board to learn which Class B species are designated for control in your area.

Dalmatian toadflax (above) is limited in distribution in Western WA, where it is a Class B Non-widespread in many counties. As with Scotch broom, Designate in regions with widespread distribution can select this species for control to meet local needs and priorities.



Scotch broom (above) is widespread in western Washington but still limited or absent in eastern Washington. The goal is to contain it where it is already abundant and to control and prevent its spread so that it does not become a problem statewide. Therefore, it is designated for control in almost every county east of the Cascades.



Non-native, invasive species that are abundant in some areas of the state, but absent or limited in other areas. The statewide goal is to "draw the line" around and contain infested regions, to keep these noxious weeds from spreading into new areas. They are designated for mandatory control in areas where they have not yet invaded or where distribution is still limited. In regions where a Class B species is already abundant, control is decided at the local level, with containment as the primary goal.

Class B Weeds

County Contacts

Snohomish	(425) 388-7548
Spokane	(509) 477-5777
Stevens	(509) 684-7590
Thurston	(360) 786-5576
Wahkiakum	(360) 795-3852
Walla Walla	(509)-524-2688
Whatcom	(360) 778-6234
Whitman	(509) 397-6261
Yakima	(509) 574-2180

To learn more about noxious weeds and noxious weed control in Washington State, please contact:

WA State Noxious Weed Control Board
P.O. Box 42560
Olympia, WA 98504-2560
(360)-725-5764

Email: noxiousweeds@agr.wa.gov
Website: <http://www.nwcb.wa.gov>

Or

WA State Department of Agriculture
21 North First Avenue #103
Yakima, WA 98902
(509) 249-6973



Cover photos of South American spongeplant infestation and plant with female flower by Jenifer Parsons, WA Department of Ecology

Eradication of Class A plants is required by law.

Garlic mustard (above) is highly invasive in many parts of the country where it completely dominates forest floors. Fortunately, its distribution is still limited in Washington. As a Class A noxious weed, it is aggressively being targeted for eradication so it does not become widespread in our forests.



Non-native, invasive plants that are limited in their distribution, and the goal is to prevent them from gaining a foothold in Washington. By law, all Class A noxious weed plants must be eradicated (eliminated).

Class A Weeds

- Prevent small infestations from expanding by eradicating them when they are first detected
- Restrict already established weed populations to regions of the state where they occur and prevent their movement to un-infested areas
- Allow flexibility of weed control at the local level for weeds that are already widespread.

To help protect the State's resources and economy, the Washington State Noxious Weed Control Board adopts a State Noxious Weed List each year (WAC 16-750). This list groups weeds into three major classes – A, B, and C – based on the distribution of each species and the seriousness of the threat they pose to Washington State. This classification system is designed to:

2020 Washington State Noxious Weed List



South American spongeplant, *Limnobium laevigatum*, is a new Class A noxious weed for 2020. Eradication in Washington is now required of this floating, aquatic perennial plant.

List arranged alphabetically by:
COMMON NAME



Noxious Weeds are non-native plants introduced to Washington State that can be highly destructive, competitive, and difficult to control. These plants invade our croplands, rangeland, forests, parks, rivers, lakes, wetlands, and estuaries causing both ecological and economical damage that affects us all.

Noxious weeds can:

- Lower crop yields
- Reduce forage quality
- Destroy plant and animal habitat
- Displace native plants
- Clog waterways
- Reduce recreational opportunities (e.g., fishing, hunting, swimming and hiking)
- Decrease land values
- Increase erosion and wildfire risk
- And some are toxic to humans and livestock

2020 Washington State Noxious Weed List

Class A Noxious Weeds - Eradication required

common crupina	<i>Crupina vulgaris</i>
cordgrass, common	<i>Spartina anglica</i>
cordgrass, dense-flowered	<i>Spartina densiflora</i>
cordgrass, saltmeadow	<i>Spartina patens</i>
cordgrass, smooth	<i>Spartina alterniflora</i>
dyer's woad	<i>Isatis tinctoria</i>
eggleaf spurge	<i>Euphorbia oblongata</i>
false brome	<i>Brachypodium sylvaticum</i>
floating primrose-willow	<i>Ludwigia peploides</i>
flowering rush	<i>Butomus umbellatus</i>
French broom	<i>Genista monspessulana</i>
garlic mustard	<i>Alliaria petiolata</i>
giant hogweed	<i>Heracleum mantegazzianum</i>
goatsrue	<i>Galega officinalis</i>
hydrilla	<i>Hydrilla verticillata</i>
Johnsongrass	<i>Sorghum halepense</i>
knapweed, bighead	<i>Centaurea macrocephala</i>
knapweed, Vochin	<i>Centaurea nigrescens</i>
kudzu	<i>Pueraria montana var. lobata</i>
meadow clary	<i>Salvia pratensis</i>
oriental clematis	<i>Clematis orientalis</i>
purple starthistle	<i>Centaurea calcitrapa</i>
reed sweetgrass	<i>Glyceria maxima</i>
ricefield bulrush	<i>Schoenoplectus mucronatus</i>
sage, clary	<i>Salvia sclarea</i>
sage, Mediterranean	<i>Salvia aethiopis</i>
silverleaf nightshade	<i>Solanum elaeagnifolium</i>
small-flowered jewelweed	<i>Impatiens parviflora</i>
South American spongeplant	<i>Limnobium laevigatum</i>
Spanish broom	<i>Spartium junceum</i>
Syrian beancaper	<i>Zygophyllum fabago</i>
Texas blueweed	<i>Helianthus ciliaris</i>
thistle, Italian	<i>Carduus pycnocephalus</i>
thistle, milk	<i>Silybum marianum</i>
thistle, slenderflower	<i>Carduus tenuiflorus</i>
variable-leaf milfoil	<i>Myriophyllum heterophyllum</i>
wild four-o'clock	<i>Mirabilis nyctaginea</i>

Class B Noxious Weeds

blueweed	<i>Echium vulgare</i>
Brazilian elodea	<i>Egeria densa</i>
bugloss, annual	<i>Anchusa arvensis</i>
bugloss, common	<i>Anchusa officinalis</i>
butterfly bush	<i>Buddleja davidii</i>
camelthorn	<i>Alhagi maurorum</i>
common fennel, (except bulbing fennel)	<i>Foeniculum vulgare</i> except <i>F. vulgare</i> var. <i>azoricum</i>)
common reed (nonnative genotypes only)	<i>Phragmites australis</i>
Dalmatian toadflax	<i>Linaria dalmatica</i> ssp. <i>dalmatica</i>
Eurasian watermilfoil	<i>Myriophyllum spicatum</i>
European coltsfoot	<i>Tussilago farfara</i>
fanwort	<i>Cabomba caroliniana</i>

Class B Noxious Weeds-Continued

gorse	<i>Ulex europaeus</i>
grass-leaved arrowhead	<i>Sagittaria graminea</i>
hairy willowherb	<i>Epilobium hirsutum</i>
hawkweed oxtongue	<i>Picris hieracioides</i>
hawkweed, orange	<i>Hieracium aurantiacum</i>
hawkweeds: All nonnative species and hybrids of the meadow subgenus	<i>Hieracium</i> , subgenus <i>Pilosella</i>
hawkweeds: All nonnative species and hybrids of the wall subgenus	<i>Hieracium</i> , subgenus <i>Hieracium</i>
herb-Robert	<i>Geranium robertianum</i>
hoary alyssum	<i>Berteroa incana</i>
houndstongue	<i>Cynoglossum officinale</i>
indigobush	<i>Amorpha fruticosa</i>
knapweed, black	<i>Centaurea nigra</i>
knapweed, brown	<i>Centaurea jacea</i>
knapweed, diffuse	<i>Centaurea diffusa</i>
knapweed, meadow	<i>Centaurea x moncktonii</i>
knapweed, Russian	<i>Rhaponticum repens</i>
knapweed, spotted	<i>Centaurea stoebe</i>
knotweed, Bohemian	<i>Polygonum x bohemicum</i>
knotweed, giant	<i>Polygonum sachalinense</i>
knotweed, Himalayan	<i>Persicaria wallichii</i>
knotweed, Japanese	<i>Polygonum cuspidatum</i>
kochia	<i>Bassia scoparia</i>
lesser celandine	<i>Ficaria verna</i>
loosestrife, garden	<i>Lysimachia vulgaris</i>
loosestrife, purple	<i>Lythrum salicaria</i>
loosestrife, wand	<i>Lythrum virgatum</i>
Malta starthistle	<i>Centaurea melitensis</i>
parrotfeather	<i>Myriophyllum aquaticum</i>
perennial pepperweed	<i>Lepidium latifolium</i>
poison hemlock	<i>Conium maculatum</i>
policeman's helmet	<i>Impatiens glandulifera</i>
puncturevine	<i>Tribulus terrestris</i>
Ravenna grass	<i>Saccharum ravennae</i>
rush skeletonweed	<i>Chondrilla juncea</i>
saltcedar	<i>Tamarix ramosissima</i>
Scotch broom	<i>Cytisus scoparius</i>
shiny geranium	<i>Geranium lucidum</i>
spurge flax	<i>Thymelaea passerina</i>
spurge laurel	<i>Daphne laureola</i>
spurge, leafy	<i>Euphorbia virgata</i>
spurge, myrtle	<i>Euphorbia myrsinites</i>
sulfur cinquefoil	<i>Potentilla recta</i>
tansy ragwort	<i>Jacobaea vulgaris</i>
thistle, musk	<i>Carduus nutans</i>
thistle, plumeless	<i>Carduus acanthoides</i>
thistle, Scotch	<i>Onopordum acanthium</i>
velvetleaf	<i>Abutilon theophrasti</i>
water primrose	<i>Ludwigia hexapetala</i>
white bryony	<i>Bryonia alba</i>
wild chervil	<i>Anthriscus sylvestris</i>
yellow archangel	<i>Lamiastrum galeobdolon</i>
yellow floatingheart	<i>Nymphoides peltata</i>
yellow nutsedge	<i>Cyperus esculentus</i>
yellow starthistle	<i>Centaurea solstitialis</i>

Class C Noxious Weeds

absinth wormwood	<i>Artemisia absinthium</i>
Austrian fieldcress	<i>Rorippa austriaca</i>
babysbreath	<i>Gypsophila paniculata</i>
black henbane	<i>Hyoscyamus niger</i>
blackgrass	<i>Alopecurus myosuroides</i>
buffalobur	<i>Solanum rostratum</i>
cereal rye	<i>Secale cereale</i>
common barberry	<i>Berberis vulgaris</i>
common catsear	<i>Hypochaeris radicata</i>
common groundsel	<i>Senecio vulgaris</i>
common St. Johnswort	<i>Hypericum perforatum</i>
common tansy	<i>Tanacetum vulgare</i>
common teasel	<i>Dipsacus fullonum</i>
curlyleaf pondweed	<i>Potamogeton crispus</i>
English hawthorn	<i>Crataegus monogyna</i>
English ivy - four cultivars only (includes Atlantic ivy)	<i>Hedera helix</i> 'Baltica', 'Pittsburgh', and 'Star'; <i>H. hibernica</i> 'Hibernica'
Eurasian watermilfoil hybrid	<i>Myriophyllum spicatum</i> x <i>M. sibiricum</i>
evergreen blackberry	<i>Rubus laciniatus</i>
field bindweed	<i>Convolvulus arvensis</i>
fragrant waterlily	<i>Nymphaea odorata</i>
hairy whitetop	<i>Lepidium appelianum</i>
Himalayan blackberry	<i>Rubus armeniacus</i>
hoary cress	<i>Lepidium draba</i>
Italian arum	<i>Arum italicum</i>
Japanese eelgrass	<i>Zostera japonica</i>
jointed goatgrass	<i>Aegilops cylindrica</i>
jubata grass	<i>Cortaderia jubata</i>
lawnweed	<i>Soliva sessilis</i>
longspine sandbur	<i>Cenchrus longispinus</i>
medusahead	<i>Taeniatherum caput-medusae</i>
nonnative cattail species and hybrids (reminder, does not include the native common cattail, <i>Typha latifolia</i>)	<i>Typha species</i>
old man's beard	<i>Clematis vitalba</i>
oxeye daisy	<i>Leucanthemum vulgare</i>
pampas grass	<i>Cortaderia selloana</i>
perennial sowthistle	<i>Sonchus arvensis</i> ssp. <i>arvensis</i>
reed canarygrass	<i>Phalaris arundinacea</i>
Russian olive	<i>Elaeagnus angustifolia</i>
scentless mayweed	<i>Matricaria perforata</i>
smoothseed alfalfa dodder	<i>Cuscuta approximata</i>
spikeweed	<i>Centromadia pungens</i>
spiny cocklebur	<i>Xanthium spinosum</i>
spotted jewelweed	<i>Impatiens capensis</i>
Swainsonpea	<i>Sphaerophysa salsula</i>
thistle, bull	<i>Cirsium vulgare</i>
thistle, Canada	<i>Cirsium arvense</i>
tree-of-heaven	<i>Ailanthus altissima</i>
ventenata	<i>Ventenata dubia</i>
white cockle	<i>Silene latifolia</i> ssp. <i>alba</i>
wild carrot (except where commercially grown)	<i>Daucus carota</i>
yellow flag iris	<i>Iris pseudacorus</i>
yellow toadflax	<i>Linaria vulgaris</i>